

Partial extraction therapy

A graftless solution for adjacent implants in the aesthetic zone

DR ALI TUNKIWALA , DR UDATTA KHER , DANESH VAZIFDAR , DR MAHENDRANATH REDDY ,
DR JOSE EDUARDO MATE SANCHEZ DE VAL

Literature

1. Sato N, Kuwana T, Yamamoto M, Suenaga H, Anada T, Koyama S, Suzuki O, Sasaki K. Bone response to immediate loading through titanium implants with different surface roughness in rats. *K.Odontology*. 2014 Jul;102(2):249-58. doi: 10.1007/s10266-013-0107-4. Epub 2013 Apr 7.
2. Esposito M, Barausse C, Pistilli R, Jacotti M, Grandi G, Tuco L, Felice P. Immediate loading of post-extractive versus delayed placed single implants in the anterior maxilla: outcome of a pragmatic multicenter randomised controlled trial 1-year after loading. *Eur J Oral Implantol*. 2015 Winter;8(4):347-58.
3. Salama M, Ishikawa T, Salama H, Funato A, Garber D. Advantages of the root submergence technique for pontic site development in esthetic implant therapy. *Int J Periodontics Restorative Dent* 2007 Dec;27(6):521-527.
4. Hürzeler MB, Zehr O, Schupbach P, Rebele SF, Emmanouilidis N, Fickl S. The socket-shield technique: a proof-of-principle report. *J Clin Periodontol* 2010 Sep;37(9):855-862.
5. Farronato D, Santoro G, Canullo L, Botticelli D, Maiorana C, Lang NP. Establishment of the epithelial attachment and connective tissue adaptation to implants installed under the concept of “platform switching”: a histologic study in minipigs. *Clin Oral Implants Res*. 2012 Jan;23(1):90-4. doi: 10.1111/j.1600-0501.2011.02196.x. Epub 2011 Apr 15.